



**BARCELONA DUST FORECAST CENTER
(WMO Regional Specialized Meteorological Center with activity specialization on
Atmospheric Sand and Dust Forecast)**

Activity Report 2014

**Enric Terradellas (AEMET), Sara Basart (BSC-CNS), José M. Baldasano (BSC-
CNS) and Emilio Cuevas (AEMET)**

Barcelona, 11 February 2015



Barcelona Dust Forecast Center
Barcelona Supercomputing Center
Nexus II Building
Jordi Girona, 29
08034 Barcelona (Spain)

Phone: (+34) 93 413 76 12
Fax: (+34) 93 413 77 21
Website: <http://dust.aemet.es>
E-mail: dust@aemet.es

Index

Introduction	4
Dust forecasts	4
The NMMB/BSC-Dust model	5
Near-Real-Time Evaluation	6
AERONET-based evaluation scores	8
Time-averaged products	9
Training	10
Health applications	10
Staff	11
Users	11
References	11

Introduction

In May 2007, the 14th WMO Congress endorsed the launching of the **Sand and Dust Storm Warning Advisory and Assessment System** (WMO SDS-WAS, [http://www.wmo.int/pages/prog/arep/wwrp/new/Sand and Dust Storm.html](http://www.wmo.int/pages/prog/arep/wwrp/new/Sand%20and%20Dust%20Storm.html)). The SDS-WAS mission is to enhance the ability of countries to deliver timely and quality sand and dust storm forecasts, observations, information and knowledge to users through an international partnership of research and operational communities. It works as an international network of research, operational centres and users, organized through regional nodes and coordinated by the SDS-WAS Steering Committee. Two nodes are currently in operation:

- Regional Node for Northern Africa, Middle East and Europe, coordinated by a Regional Center in Barcelona, Spain, hosted by the State Meteorological agency (AEMET) and the Barcelona Supercomputing Center (BSC-CNS): <http://sds-was.aemet.es/>
- Regional Node for Asia, coordinated by a Regional Center in Beijing, China, that is hosted by the China Meteorological Administration (CMA): <http://www.sds.cma.gov.cn/>

In May 2013, in view of the demand of many national meteorological services and the good results obtained by the SDS-WAS, which prove the feasibility and the need to begin developing operational services beyond the scope of R&D, the 65th Session of the WMO Executive Council designates the consortium formed by AEMET and the BSC-CNS to create in Barcelona the first Regional Specialized Meteorological Center with activity specialization on Atmospheric Sand and Dust Forecast (RSMC-ASDF). The Center shall operationally generate and distribute predictions for Northern Africa (north of equator), Middle East and Europe. The **Barcelona Dust Forecast Center** (BDFC, <http://dust.aemet.es>) was created in February 2014 to be this RSMC-ASDF and was publicly presented on 10 June 2014.

Dust forecasts

The **Barcelona Dust Forecast Center** (BDFC) prepares regional forecast fields using the **NMMB/BSC-Dust** model continuously throughout the year on a daily basis. The model consists of a numerical weather prediction model incorporating on-line parameterizations of all the major phases of the atmospheric dust cycle. It is run at a horizontal resolution of 0.1 degrees longitude per 0.1 degrees latitude for a domain covering Northern Africa, Middle East and Europe (25°W-65°E, 0°-65°N). This domain covers the main dust source areas in Northern Africa and Middle East, as well as the main transport routes and deposition zones from the equator to the Scandinavian Peninsula.

The BDFC generates forecasts of the following set of variables:

- Dust load (kg.m⁻²)
- Dust surface concentration (µg.m⁻³)
- Dust optical depth at 550 nm (-)
- 3-hour accumulated dry and wet deposition (kg.m⁻²)
- Dust surface extinction (Mm⁻¹)

Forecasts cover the period from the starting time (12 UTC) up to a lead time of 72 hours. They are distributed in pictorial form through the Center’s web portal, where are posted 12 hours after the starting forecast time and through the **WMO Global Telecommunications System (GTS)**.

The headers of the GTS bulletins are as follows:

PZBcnn LEMM dd1200, where:

- c=**A** denotes a 0-hour forecast, c=**B** a 3-hour forecast, ..., c=**Y** a 72-hour forecast
- nn=**01** denotes dust surface concentration, nn=**02** dust optical depth, nn=**03** dust dry deposition, nn=**04** dust wet deposition and nn=**05** dust load.
- dd indicates the day

The model outputs are stored, both in netcdf and in **WMO FM 92 GRIB Edition 2** formats. For GRIB files, version 13.0.0 of the code tables is used:

- Code and flag tables v. 13.0.0
- Local tables v. 1.0.0 (see Table 1). These tables have been built by the Barcelona Dust Forecast Center to complement the WMO Master Tables in order to enable encoding/decoding dust-related products.

Number	Parameter	Units
248	Dust surface extinction at 550 nm	Mm ⁻¹
249	Dust load	g.m ⁻²
250	3-hour accumulated dust dry deposition	mg.m ⁻²
251	3-hour accumulated dust wet deposition	mg.m ⁻²
252	Dust surface concentration	µg.m ⁻³
253	Dust optical depth at 550 nm	-

Table 1. Code Table 4.2 - Parameter number by product discipline and parameter category. **Product discipline 0** - Meteorological products, parameter category 13: aerosols

The NMMB/BSC-Dust model

The NMMB/BSC-Dust (Pérez et al., 2011; Haustein et al., 2012) is an online multi-scale atmospheric dust model designed and developed at the BSC-CNS) in collaboration with the **National Centers for Environmental Prediction (NCEP)**, the **Goddard Institute for Space Studies** and the **International Research Institute for Climate and Society (IRI)**. The dust model is fully embedded into the **Non-hydrostatic Multiscale Model** on B-grid (NMMB) model developed at NCEP (Janjic, 2005; Janjic and Black, 2007; Janjic et al., 2011) and is intended to provide short to medium-range dust forecasts for both regional and global domains.

The NMMB/BSC-Dust model solves the mass balance equation for dust taking into account the following processes:

- Dust generation and uplift by surface wind and turbulence. The emission scheme considers saltation and sandblasting processes (White, 1979; Marticorena and Bergametti, 1995) and the presence of a viscous sublayer between the surface

and the lowest model layer (Janjic, 1994). The size distribution of soil particles is based on the soil textures of the STATSGO-FAO soil map. Then, the dust vertical flux is distributed over eight size bins. For the source function, the model uses the topographic preferential source approach described by Ginoux et al. (2001) and climatological vegetation data from the National Environmental Satellite, Data, and Information Service (NESDIS).

- Soil wetness effects on dust production (Fécan, 1999).
- Horizontal and vertical advection (Janjic et al., 2009).
- Horizontal diffusion and vertical transport by turbulence and convection (Janjic et al., 2009).
- Dry deposition and gravitational settling (Zhang et al., 2001).
- Wet removal, which includes in-cloud and below-cloud scavenging from convective and stratiform clouds (Betts, 1986; Janjic, 1994; Ferrier et al., 2002).
- Furthermore, in order to take into account the radiative effects of mineral dust, the rapid radiative transfer model (RRTM) (Mlawer et al., 1997) has been implemented.

Near-Real-Time Evaluation

The Near-Real-Time (NRT) model evaluation is an assessment of how the forecast behaves relative to observations that are NRT available, allowing the modelling group and the end users to have a quick overview of the quality of the forecast. Forecasts of dust optical depth (DOD) are compared with total aerosol optical depth (AOD) provided by the AErosol RObotic NETwork (AERONET, <http://aeronet.gsfc.nasa.gov/>) for selected dust-prone stations located in Northern Africa, Middle East and Southern Europe. At present, 40 AERONET stations are used. They are listed in Table 2.

Site name	Longitude	Latitude	Country	Region
Avignon	4.88°E	43.93°N	France	Mediterranean
Banizoumbou	2.66°E	13.54°N	Niger	Sahel/Sahara
Barcelona	2.12°E	41.39°N	Spain	Mediterranean
Cabo_da_Roca	9.50°W	38.78°N	Portugal	Mediterranean
Caceres	6.34°W	39.48°N	Spain	Mediterranean
Cairo_EMA_2	31.29°E	30.08°N	Egypt	Mediterranean
Capo_Verde	22.93°W	16.73°N	Cabo Verde	Sahel/Sahara
CUT-TEPAK	33.04°E	34.67°N	Cyprus	Mediterranean
Dakar	16.96°W	14.39°N	Sénégal	Sahel/Sahara
Eilat	34.92°E	29.50°N	Israel	Mediterranean
Ersa	9.36°E	43.00°N	France	Mediterranean
ETNA	15.02°E	37.61°N	Italy	Mediterranean
Evora	7.91°W	38.57°N	Portugal	Mediterranean
FORTH_CRETE	25.28°E	35.33°N	Greece	Mediterranean
Granada	3.60°W	37.16°N	Spain	Mediterranean
IASBS	48.51°E	36.70°N	Iran	Middle East
IER_Cinzana	5.93°W	13.28°N	Mali	Sahel/Sahara
Ilorin	4.34 °E	8.32°N	Nigeria	Sahel/Sahara
IMAA_Potenza	15.72°E	40.60°N	Italy	Mediterranean
IMS-METU-ERDEMLI	34.26°E	36.56°N	Turkey	Mediterranean
KAUST_Campus	39.10 °E	22.30 °N	Saudi Arabia	Middle East
Kuwait University	47.97 °E	29.33 °N	Kuwait	Middle East
Lampedusa	12.63°E	35.52°N	Italy	Mediterranean
Lecce_University	18.11°E	40.33°N	Italy	Mediterranean
Oujda	1.90°W	34.65°N	Morocco	Mediterranean
Ouarzazate	6.91°W	30.93°N	Morocco	Sahel/Sahara
Palma_de_Mallorca	2.62°E	39.55°N	Spain	Mediterranean
Porquerolles	6.16°E	43.00°N	France	Mediterranean
Rome_Tor_Vergata	12.65°E	41.84°N	Italy	Mediterranean
Saada	8.16°W	31.62°N	Morocco	Sahel/Sahara
Santa_Cruz_Tenerife	16.25°W	28.47°N	Spain	Sahel/Sahara
SEDE_BOKER	34.78°E	30.85°N	Israel	Mediterranean
Seysses	1.26°E	43.50°N	France	Mediterranean
Solar_Village	46.40°E	24.91°N	S. Arabia	Middle East
Tabernas_PSA-DLR	2.36°W	37.09°N	Spain	Mediterranean
Tamanrasset_INM	5.53°E	22.79°N	Algeria	Sahel/Sahara
Villefranche	7.33°E	43.68°N	France	Mediterranean
Xanthi	24.92°E	41.15°N	Greece	Mediterranean
Zinder_Airport	8.99°E	13.78°N	Niger	Sahel/Sahara
Zouerate-Fennec	12.48°W	22.75°N	Mauritania	Sahel/Sahara

Table 2. AERONET stations, where the evaluation has been implemented.

The AERONET program is a federation of ground-based remote sensing aerosol networks established by the U. S. National Aeronautics and Space Administration

(NASA) and the French network PHOTONS managed by the Lille 1 University, the National Centre for Space Studies (CNES) and the National Centre for Scientific Research (CNRS). The AERONET network is greatly expanded by collaborators from national agencies, institutes, universities and other partners. The program provides a long-term, continuous and readily accessible public domain database of aerosol optical, microphysical and radiative properties for aerosol research and characterization, validation of satellite retrievals, and synergism with other databases. The network imposes standardization of instruments, calibration, processing and distribution.

Version 2 Level 1.5 of AERONET products are used for the NRT evaluation. Level 1.5 data are automatically cloud screened but not have final calibration applied. However, differences between Level 1.5 and Level 2 are normally quite small if the instrument has not suffered dramatic changes between the pre and the post calibration. Anyway, these differences are normally within the error of the model predictions. Since AERONET sun photometers do not yield AOD at 550 nm, this variable is calculated from AOD at 440, 675 and 870 nm and the Ångström exponent 440-870 using the Ångström law.

AERONET-based evaluation scores

Besides the NRT evaluation, a system to evaluate the performance of the model has been set. The system yields, on a monthly, seasonal and annual basis, evaluation scores computed from the comparison of the simulated DOD and the AERONET retrievals of AOD used in the NRT evaluation. Rather than time-interpolated, retrievals are assigned to the nearest multiple-of-3 hour. In case more than one value is assigned to the same hour, only the closest-in-time is considered.

The evaluation system is applied to instantaneous forecast values of DOD ranging from the initial day (D) at 15:00 UTC to the following day (D+1) at 12:00 UTC, that is for lead times ranging from 3 to 24 hours.

To minimize the sources of error, it is intended to restrict the comparison to situations in which mineral dust is the dominant aerosol type. Threshold discrimination is made by discarding observations with an Ångström exponent 440-870 higher than 0.6. However, other particles are always present (anthropogenic aerosol, products from biomass burning, etc.) and therefore a negative bias can be expected.

The scores are computed for each AERONET site, for 3 sub-regions (Sahel/Sahara, Middle East and Mediterranean) as well as globally considering all sites (Table 2). It should be noted that scores for individual stations can be little significant for being calculated from a small number of data.

The metrics that are used to quantify the mean departure between modelled (c_i) and observed (o_i) quantities are the mean bias error (BE), the root mean square error (RMSE), the correlation coefficient (r) and the fractional gross error (FGE). They are presented in the Table 3, where n denotes the number of data.

Statistic Parameter	Formula	Range	Perfect score
Mean Bias Error (BE)	$BE = \frac{1}{n} \sum_{i=1}^n (c_i - o_i)$	$-\infty$ to $+\infty$	0
Root Mean Square Error (RMSE)	$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (c_i - o_i)^2}$	0 to $+\infty$	0
Correlation coefficient (r)	$r = \frac{\sum_{i=1}^n (c_i - \bar{c}) \cdot (o_i - \bar{o})}{\sqrt{\sum_{i=1}^n (c_i - \bar{c})^2} \cdot \sqrt{\sum_{i=1}^n (o_i - \bar{o})^2}}$	-1 to 1	1
Fractional Gross Error (FGE)	$FGE = \frac{2}{n} \sum_{i=1}^n \left \frac{c_i - o_i}{c_i + o_i} \right $	0 to 2	0

Table 3. Metrics used in the forecast evaluation.

Some considerations about these metrics are the following:

- The **mean bias error** (BE) captures the average deviations between two datasets. It has the same units as the variable. Values near 0 are the best, negative values indicate underestimation and positive values indicate overestimation.
- The **root mean square error** (RMSE) combines the spread of individual errors. It is strongly dominated by the largest values, due to the squaring operation. Especially in cases where prominent outliers occur, the usefulness of RMSE is questionable and the interpretation becomes more difficult.
- The **correlation coefficient** (r) indicates the extent to which patterns in the model match those in the observations.
- The **fractional gross error** (FGE) is a measure of model error, ranging between 0 and 2 and behaves symmetrically with respect to under- and overestimation, without over-emphasizing outliers.

The results for the whole period (starting 9 February) are summarized in Table 4.

Region / sub-region	BE	RMSE	r	FGE	n
Sahel /Sahara	-0.12	0.38	0.51	0.75	7427
Middle East	-0.10	0.27	0.39	0.64	112
Mediterranean	-0.19	0.30	0.46	1.34	4623
Total	-0.15	0.35	0.52	0.98	12162

Table 4. Annual evaluation scores (2014)

Time-averaged products

Decadal (10 days) and monthly averages of dust surface concentration are computed for long-term monitoring. Seasonal and annual averages are expected to be implemented in the near future.

The **Daily average** is computed as the mean value of the forecasts with lead times ranging from 0 to 21 hours. It means that the daily average for a specific day is an estimation of the mean value predicted over the period from 12 UTC of that day to 9 UTC of the following day. The **monthly average** is computed as the mean value of the daily averages from the first to the last day of the concerned month.

Decadal averages are not published on the web portal, but are available on demand. In particular, they are delivered to the African Center of Meteorological applications for Development (ACMAD; <http://www.acmad.net/new/>) that uses them in health-oriented programs.

Training

The 4th Training Course on WMO SDS-WAS products (satellite and ground observation and modelling of atmospheric dust) was held 17-20 November 2014 in Casablanca, Morocco. The event has been organized by the **Barcelona Dust Forecast Center**, the **Regional Center for Northern Africa, Middle East and Europe of the WMO SDS-WAS** and the **National Meteorology Direction (DMN)** of Morocco and financed by AEMET and the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT).

The event was targeted to operational meteorologists and early career scientists (advanced students, PhD candidates and postdoctoral researchers) with interest on the Earth system sciences. Participants from Algeria, Egypt, The Gambia, Iran, Morocco, Niger, Togo and ACMAD attended a series of lectures and practical exercises. In particular, they became familiar with the products distributed by the BDFC and learnt the most efficient way to introduce them in their daily work.

Health applications

The activities carried out by the Barcelona Dust Forecast Center were presented at the World Health Organization's Regional Consultation Meeting on Air Quality and Health for the Eastern Mediterranean Region held at Amman, Jordan, 10-11 December 2014 and attended by public health and environment managers from the region.

Staff

Enric Terradellas, technical director

Sara Basart, research and operations

Francesco Benincasa and **Kim Serradell**, technical support

José M. Baldasano and **Emilio Cuevas**, scientific advisers

Users

The BDFC makes a quarterly monitoring of the accesses to its web portal. The results (Table 5) show a steady increase in the number of users, sessions and page views.

Season	Users	Sessions	Page views
Spring (Mar-May)	422	878	2521
Winter (Jun-Aug)	1132	2449	6134
Autumn (Sep-Nov)	1909	3175	6901

Table 5. Access to the web portal. Quarterly review (2014)

References

Betts, A. K. and Miller, M. J.: A new convective adjustment scheme. Part 2: Single column tests using GATE wave, BOMEX, ATEX and arctic air-mass data sets, *Q. J. Roy. Meteor. Soc.*, 112, 693–709, doi:10.1002/qj.49711247308, 1986.

Fécan, F., Marticorena, B., and Bergametti, G.. Parameterization of the increase of the aeolian erosion threshold wind friction due to soil moisture for semi arid areas, *Ann. Geophys.*, 17, 149-157, 1999.

Ferrier, B. S., Jin, Y., Lin, Y., Black, T., Rogers, E., and DiMego, G.: Implementation of a new grid-scale cloud and precipitation scheme in the NCEP Eta Model, in: Proc. 15th Conf. on Numerical Weather Prediction, 12–16 August 2002, San Antonio, TX, Amer. Meteor. Soc., 280–283, 2002.

Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S.-J.: Sources and distributions of dust aerosols simulated with the GOCART model, *J. Geophys. Res.*, 106, 20255-20274 10.1029/2000JD000053 2001.

Haustein, K., Pérez, C., Baldasano, J. M., Jorba, O., Basart, S., Miller, R. L., Janjic, Z., Black, T., Nickovic, S., Todd, M. C., Washington, R., Müller, D., Tesche, M., Weinzierl, B., Esselborn, M. and Schladitz, A.: Atmospheric dust modeling from meso to global scales with the online NMMB/BSC-Dust model–Part 2: Experimental campaigns in Northern Africa, *Atmos. Chem. Phys.*, 12, 2933–2958, doi:10.5194/acp-12-2933-2012, 2012.

Janjic, Z. I.: The Step-mountain Eta Coordinate Model: Further developments of the convection, viscous sublayer and turbulence closure schemes, *Mon. Weather Rev.*, 122, 927– 945, 1994.

Janjic, Z. I.: A unified model approach from meso to global scales. *Geophys. Res. Abstracts*, 7, SRef{ID: 1607{7962/gra/EGU05}{A{05 582, 2005.

Janjic, Z. I. and Black, T: A unified model approach from meso to global scales. *Geophys. Res. Abstracts*, 7, SRef{ID: 1607{7962/gra/EGU2007}{A{05 025, 2007.

Janjic, Z., Huang, H., and Lu, S.: A unified atmospheric model suitable for studying transport of mineral aerosols from meso to global scales, *IOP C. Ser. Earth Env.*, 7, 012011, 2009.

Janjic, Z., Janjic, T., and Vasic, R.: A Class of conservative fourth order advection schemes and impact of enhanced formal accuracy on extended range forecasts, *Mon. Weather Rev.*, 0, null, doi:10.1175/2010MWR3448.1, 2011.

Marticorena, B., and Bergametti, G.: Modeling the atmospheric dust cycle: 1. Design of a soil-derived dust emission scheme, *J. Geophys. Res.*, 100, 6415-16430, 1995.

Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.*, 102, 16663–16682, 1997.

Pérez, C., Haustein, K., Janjic, Z., Jorba, O., Huneeus, N., Baldasano, J.M., Black, T., Basart, S., Nickovic, S., Miller, R.L., Perlwitz, J., Schulz, M. and Thomson, M.: An online mineral dust aerosol model for meso to global scales: Model description, annual simulations and evaluation, *Atmos. Chem. Phys.*, 11, 13001-13027, doi: 10.5194/acp-11-13001-2011, 2011.

White, B. R.: Soil transport by winds on Mars. *J. Geophys. Res.* 84, 4643-4651, 1979.

Zhang, L., Gong, S., Padro, J., and Barrie, L.: A size-segregated particle dry deposition scheme for an atmospheric aerosol module, *Atmos. Environ.*, 35, 549-560, 2001.